Reply to Office Action of January 11, 2010

AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A π -conjugated aromatic ring-containing compound, characterized by being represented by the formula (1)

[Chemical Formula 1]

$$R^{1} \xrightarrow{A} \xrightarrow{R^{2}} \left(\begin{array}{c} \\ \\ \\ \\ \\ \end{array} \right) \xrightarrow{R^{3}} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{A_{1}} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{1}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{1}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{1}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{A_{2}}$$

{wherein R¹ represents a hydrogen atom, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or chlorine atom, or a group represented by the following formula (2) or a group represented by the following formula (3)

[Chemical Formula 2]

$$E = \begin{array}{c} R^7 \\ OH \\ (2) \end{array}$$

[wherein E represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, an imidazolyl group or a thienyl group (provided that said phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom),

Application No.: 10/591,950 Reply dated July 8, 2010

Reply to Office Action of January 11, 2010

R⁷ and R⁸ independently represent an alkyl group having 1 to 10 carbon atoms],

 R^2 , R^3 , R^4 , and R^5 independently represent a hydrogen atom or an alkyl group having 1 to 10 carbon atoms.

R⁶ represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a pyridyl group, a pyridyl group, a pyridyl group, an imidazolyl group, a thienyl group (provided that the phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom), a group represented by the afore-indicated formula (2) or a group represented by the afore-indicated formula (3),

A and D independently represent a naphthalene ring, an anthracene ring, a phenanthrene group, a phenarene ring, a fluorene ring, a triphenylene ring, a pyrene ring, a perylene ring, a pyridine ring, a thiophene ring, a benzothiadiazole ring, a thieno[3,4-b]pyrazine ring, a furo[3,4-b]pyrazine ring or a 6H-pyrrolo[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom),

a₁, a₂, and a₃ are independently 0 or 1, and
n₁ and n₂ are independently an integer of 1 to 5}.

2. (Currently amended) A π -conjugated aromatic ring-containing compound, characterized by being represented by the formula (4)

Docket No.: 0171-1307PUS1

Page 3 of 21

Docket No.: 0171-1307PUS1 Page 4 of 21

Application No.: 10/591,950
Reply dated July 8, 2010
Reply to Office Action of January 11

Reply to Office Action of January 11, 2010

[Chemical Formula 3]

$$R^{6} = \left[\begin{array}{c} Y^{1} = & \\ & \\ & \end{array} \right] \xrightarrow{R^{2}} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} \\ \\ \\ \end{array} \right) \xrightarrow{d} \left(\begin{array}{c}$$

[wherein R², R³, R⁴, and R⁵ independently represent a hydrogen atom or an alkyl group having 1 to 10 carbon atoms,

R⁶ represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a pyridyl group, a pyridyl group, a pyrazolyl group, an imidazolyl group, a thienyl group (provided that the phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom), a group represented by the formula (2) or a group represented by the formula (3),

[Chemical Formula 4]

$$E \xrightarrow{\circ} R^{8} \xrightarrow{R^{7}} OH$$
(2) (3)

[wherein E represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a pyridyl group, a pyridyl group, a pyrazinyl group, a furanyl group, a pyrrolyl group, a pyrazolyl group, an imidazolyl group or a thienyl group (provided that said phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a

Application No.: 10/591,950 Docket No.: 0171-1307PUS1
Reply dated July 8, 2010 Page 5 of 21

Reply to Office Action of January 11, 2010

dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom),

R⁷ and R⁸ independently represent an alkyl group having 1 to 10 carbon atoms],

Z¹, Y¹, and Y² independently represent a benzene ring, a paraphenylene group, a naphthalene ring, an anthracene ring, a phenanthrene group, a phenarene ring, a fluorene ring, a triphenylene ring, a pyrene ring, a perylene ring, a pyridine ring, a pyrimidine ring, a pyridazine ring, a pyrazine ring, a furan ring, a pyrrole ring, a pyrazole ring, an imidazole ring, a thiophene ring, a benzothiadiazole ring, a thieno[3,4-b]pyrazine ring, a furo[3,4-b]pyrazine ring or a 6H-pyrrolo[3,4-b]pyrazine ring (provided that these rings and group may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom), R⁹ represents a single bond, -O-, -S-, -S(O)-, -S(O₂)-, -C(O)O-, -OC(O)-, -C(S)O-, -OC(S)-, -C(O)NH-, -NHC(O)-, -C(S)NH-, -NHC(S)-, -NH- or a divalent saturated or unsaturated hydrocarbon group that has 1 to 8 carbon atoms and may be branched,

b₁ and b₂ are independently 0 or 1,
c is an integer of 0 to 3, and
m₁ and m₂ are independently an integer of 1 to 5].

3. (Previously Presented) A π -conjugated aromatic ring-containing compound, characterized by being represented by the formula (5)

Application No.: 10/591,950 Docket No.: 0171-1307PUS1
Reply dated July 8, 2010 Page 6 of 21

Reply to Office Action of January 11, 2010

[Chemical Formula 5]

$$R^{10}$$

$$R^{10}$$

$$R^{10}$$

$$R^{11}$$

$$R^{2}$$

$$R^{3}$$

$$R^{2}$$

$$R^{3}$$

$$R^{4}$$

$$R^{5}$$

$$R^{4}$$

$$R^{5}$$

$$R^{4}$$

$$R^{6}$$

$$R^{5}$$

$$R^{6}$$

$$R^{2}$$

$$R^{2}$$

$$R^{4}$$

$$R^{6}$$

$$R^{5}$$

$$R^{6}$$

$$R^{2}$$

$$R^{4}$$

$$R^{6}$$

$$R^{6}$$

$$R^{2}$$

$$R^{3}$$

$$R^{4}$$

$$R^{6}$$

$$R^{5}$$

$$R^{4}$$

$$R^{6}$$

$$R^{2}$$

$$R^{3}$$

$$R^{4}$$

$$R^{6}$$

$$R^{2}$$

$$R^{4}$$

$$R^{6}$$

$$R^{2}$$

$$R^{3}$$

$$R^{4}$$

$$R^{6}$$

$$R^{2}$$

$$R^{3}$$

$$R^{4}$$

$$R^{6}$$

$$R^{2}$$

$$R^{3}$$

{wherein R², R³, R⁴, and R⁵ independently represent a hydrogen atom or an alkyl group having 1 to 10 carbon atoms,

R⁶ represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a hienyl group (provided that the phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom), a group represented by the formula (2) or a group represented by the formula (3),

Application No.: 10/591,950 Reply dated July 8, 2010

Docket No.: 0171-1307PUS1 Page 7 of 21 Reply to Office Action of January 11, 2010

[Chemical Formula 6]

[wherein E represents a hydrogen atom, a substituted silvl group, a phenyl group, a naphthyl group, a pyridyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazinyl group, a furanyl group, a pyrrolyl group, a pyrazolyl group, an imidazolyl group or a thienyl group (provided that said phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms, or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine or chlorine atom),

R⁷ and R⁸ independently represent an alkyl group having 1 to 10 carbon atoms],

 R^{10} and R^{11} independently represent a hydrogen atom or an alkyl group having 1 to 10 carbon atoms.

Y³ to Y⁵ independently represent a benzene ring, a naphthalene ring, an anthracene ring, a phenanthrene group, a phenarene ring, a fluorene ring, a triphenylene ring, a pyrene ring, a perylene ring, a pyridine ring, a pyrimidine ring, a pyridazine ring, a pyrazine ring, a furan ring, a pyrrole ring, a pyrazole ring, an imidazole ring, a thiophene ring, a benzothiadiazole ring, a thieno[3,4-b]pyrazine ring, a furo[3,4-b]pyrazine ring or a 6H-pyrrolo[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom),

Z² represents a trivalent aryl group, a group represented by the following formula (6) or a group represented by the following formula (7)

Reply to Office Action of January 11, 2010

Docket No.: 0171-1307PUS1 Page 8 of 21

[Chemical Formula 7]

d₁ to d₃ are independently 0 or 1, and

 k_1 to k_3 are independently an integer of 1 to 5}.

4. (Original) The π -conjugated aromatic ring-containing compound according to Claim 1, characterized in that said R^1 is a hydrogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, a methoxy group, a propoxy group, a methyl group, a trifluoromethyl group, a group represented by the following formula (8) or a group represented by the following formula (9)

[Chemical Formula 8]

[wherein E' represents a hydrogen atom, a trimethylsilyl group, a tri-i-propylsilyl group, a phenyl group, a pyridyl group, a thienyl group (provided that the phenyl group, pyrdyl group or thienyl group may be optionally substituted with a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, a methoxy group, an n-propoxy group, a methyl group or a trifluoromethyl group)].

5. (Original) The π -conjugated aromatic ring-containing compound according to any one of Claims 1 to 4, characterized in that said R^2 , R^3 , R^4 , and R^5 independently represent a hydrogen atom, a methyl group, an ethyl group or an n-propyl group.

Application No.: 10/591,950 Reply dated July 8, 2010 Reply to Office Action of January 11, 2010

Docket No.: 0171-1307PUS1 Page 9 of 21

6. (Previously Presented) The π -conjugated aromatic ring-containing compound according to Claim 1, characterized in that said R⁶ is a hydrogen atom, a trimethylsilyl group, a tri-i-propylsilyl group, a phenyl group, a pyridyl group, a thienyl group (provided that said phenyl group, pyridyl group or thienyl group may be optionally substituted with a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, a methoxy group, an n-propoxy group, a methyl group or a trifluoromethyl group), a group represented by the following formula (8) or a group represented by the following formula (9),

[Chemical Formula 9]

$$E'$$
 CH_3 OH OH (9)

[wherein E' represents a hydrogen atom, a trimethylsilyl group, a tri-i-propylsilyl group, a phenyl group, a pyridyl group, a thienyl group (provided that the phenyl group, pyrdyl group or thienyl group may be optionally substituted with a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, a methoxy group, an n-propoxy group, a methyl group or a trifluoromethyl group)].

7. (Currently Amended) The π -conjugated aromatic ring-containing compound according to Claim 1, characterized in that said A and D independently represent a pyridine ring, a pyrimidine ring, a pyridazine ring, a pyrazine ring, a furan ring, a pyrrole ring, a pyrazole ring, an imidazole ring, a thiophene ring, a benzothiadiazole ring, a thieno[3,4-b]pyrazine ring, a furo[3,4-b]pyrazine ring or a 6H-pyrrolo[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom).

Application No.: 10/591.950 Docket No.: 0171-1307PUS1 Page 10 of 21

Reply dated July 8, 2010

Reply to Office Action of January 11, 2010

8. (Currently Amended) The π -conjugated aromatic ring-containing compound according to Claim 7, characterized in that said A and D independently represent-a pyridine ring, a pyridazine ring, a thiophene ring, a benzothiadiazole ring or a thieno[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a cyano group, a methoxy group, an n-propoxy group, a methyl group or a trifluoromethyl group).

- 9. (Previously Presented) The π -conjugated aromatic ring-containing compound according to Claim 2, characterized in that said Z¹, Y¹, and Y² independently represent a benzene ring, a naphthalene ring, an anthracene ring, a pyridine ring, a pyridazine ring, a thiophene ring, a pyrrole ring, a benzothiadiazole ring or a thieno[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a cyano group, a methoxy group, an npropoxy group, a methyl group or a trifluoromethyl group), and R⁹ represents a single bond or -O-, b_1 and b_2 are both 1, and c is 0 or 1.
- 10. (Previously Presented) The π -conjugated aromatic ring-containing compound according to Claim 3, characterized in that said Y³ to Y⁵ independently represent a phenylene ring, a naphthalene ring, an anthracene ring, a pyridine ring, a pyridazine ring, a thiophene ring, a pyrrole ring, a benzothiadiazole ring or a thieno[3,4-b]pyrazine ring (provided that these rings may be optionally substituted with a phenyl group, a cyano group, a methoxy group, an npropoxy group, a methyl group or a trifluoromethyl group), Z² is a group represented by the following formula (10) or a group represented by the following formula (11), and d₁ to d₃ are all 1.

[Chemical Formula 10]

Docket No.: 0171-1307PUS1

Page 11 of 21

Application No.: 10/591,950 Reply dated July 8, 2010

Reply to Office Action of January 11, 2010

11. (Previously Presented) An organic electroluminescent device of a type which comprises an anode and a cathode, and an organic thin film layer interposed their between, characterized in that said organic thin film layer is a layer constituted to contain the π -conjugated aromatic ring-containing compound defined in Claim 1.

12. (New) A π -conjugated aromatic ring-containing compound, characterized by being represented by the formula (12)

[Chemical Formula 11]

$$\begin{array}{c|c}
R^1 & & & \\
\hline
 & & & \\
R^3 & & & \\
\hline
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\
 & & & \\
\hline
 & & & \\$$

{wherein R¹ represents a hydrogen atom, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 atoms, a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or chlorine atom, or a group represented by the following formula (2)

[Chemical Formula 12]

$$F =$$
 (2)

[wherein E represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a pyridinyl group, a pyridazinyl group, a pyrazinyl group, a furanyl group, a pyrrolyl group, a pyrazolyl group, an imidazolyl group or a thienyl group (provided that said phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl

Reply to Office Action of January 11, 2010

Docket No.: 0171-1307PUS1 Page 12 of 21

group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 atoms, a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or chlorine atom)],

R² and R³ independently represent a hydrogen atom or an alkyl group having 1 to 10 carbon atoms,

R⁶ represents a hydrogen atom, a substituted silyl group, a phenyl group, a naphthyl group, a pyridyl group, a pyrimidinyl group, a pyridazinyl group, a pyrazinyl group, a furanyl group, a pyrrolyl group, a pyrazolyl group, an imidazolyl group or a thienyl group (provided that said phenyl group, naphthyl group, pyridyl group, pyrimidinyl group, pyridazinyl group, pyrazinyl group, furanyl group, pyrrolyl group, pyrazolyl group, imidazolyl group or thienyl group may be optionally substituted with a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 atoms, a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or chlorine atom), or a group represented by the following formula (3),

[Chemical Formula 13]

$$R^8 \xrightarrow{R^7}$$
 (3)

(wherein R⁷ and R⁸ independently represent an alkyl group having 1 to 10 carbons atoms),

A and D independently represent a pyridine ring or a thiophene ring (provided that these rings may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom),

 a_1 is independently 0 or 1, and

Application No.: 10/591,950 Reply dated July 8, 2010

Reply to Office Action of January 11, 2010

Docket No.: 0171-1307PUS1 Page 13 of 21

 n_2 is independently an integer or 1 to 5}.

13. (New) The π -conjugated aromatic ring-containing compound of claim 12, wherein R¹ represents a hydrogen atom, an alkyl group having 1 to 10 carbon atoms or a group represented by the following formula (2)

[Chemical Formula 14]

$$\mathbf{F}$$
 (2)

[wherein E represents a hydrogen atom, a substituted silyl group, a thienyl group (provided that said thienyl group may be optionally substituted with a cyano group)],

R⁶ represents a hydrogen atom, a substituted silyl group, (provided that said pyridyl group may be optionally substituted with a cyano group or an alkoxy group having 1 to 3 carbon atoms), or a group represented by the following formula (3),

[Chemical Formula 13]

$$R^8 \xrightarrow{R^7}$$
 (3)

(wherein R⁷ and R⁸ represent an alkyl group having 1 to 10 carbon atoms).

- 14. (New) The π -conjugated aromatic ring-containing compound of claim 13, wherein R⁷ and R⁸ represent a methyl group.
- 15. (New) The π -conjugated aromatic ring-containing compound of claim 13, wherein Z^1 , Y^1 and Y^2 independently represent a naphthalene ring, an anthracene ring, a phenanthrene group, a phenarene ring, a fluorene ring, a triphenylene ring, a pyrene ring, a perylene ring, a pyridine ring, a pyrimidine ring, a pyridazine ring, a pyrazine ring, a furan ring, a pyrrole ring, a pyrazole ring, an imidazole ring, a thiophene ring, a benzothiadiazole ring, a thieno[3,4-b]pyrazine ring, a furo[3,4-b]pyrazine ring or a 6H-pyrrolo[3,4-b]pyrazine ring (provided that

Docket No.: 0171-1307PUS1 Application No.: 10/591,950 Page 14 of 21

Reply dated July 8, 2010

Reply to Office Action of January 11, 2010

these rings may be optionally substituted with a phenyl group, a halogen atom, a cyano group, a nitro group, a dimethylamino group, a diphenylamino group, an alkoxy group having 1 to 3 carbon atoms, an alkyl group having 1 to 10 carbon atoms or a halogenated alkyl group that has 1 to 10 carbon atoms and may be optionally substituted with a fluorine atom or a chlorine atom).